

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
6 May 2004 (06.05.2004)

PCT

(10) International Publication Number  
**WO 2004/037786 A2**

(51) International Patent Classification<sup>7</sup>: C07D 207/00

3L1 (CA). WILSON, Marie-Claire [CA/CA]; 16711  
Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA).

(21) International Application Number:

PCT/CA2003/001620

(74) Agent: OGILVY RENAULT; Attn: Murphy, Kevin.  
P., Suite 1600, 1981 McGill College Avenue, Montreal,  
Québec H3A 2Y3 (CA).

(22) International Filing Date: 23 October 2003 (23.10.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/421,402

25 October 2002 (25.10.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,  
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*):  
MERCK FROSST CANADA & CO. [CA/CA]; 16711  
Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BILLOT, Xavier  
[FR/CA]; 16711 Trans-Canada Highway, Kirkland,  
Québec H9H 3L1 (CA). BEUNARD, Jean-Luc [FR/CA];  
16711 Trans-Canada Highway, Kirkland, Québec H9H  
3L1 (CA). HAN, Yongxin [CA/CA]; 16711 Trans-Canada  
Highway, Kirkland, Québec H9H 3L1 (CA). YOUNG,  
Robert, N. [CA/CA]; 16711 Trans-Canada Highway,  
Kirkland, Québec H9H 3L1 (CA). COLUCCI, John  
[CA/CA]; 16711 Trans-Canada Highway, Kirkland,  
Québec H9H 3L1 (CA). GIRARD, Mario [CA/CA];  
16711 Trans-Canada Highway, Kirkland, Québec H9H

Published:

— without international search report and to be republished  
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: EP<sub>4</sub> RECEPTOR AGONISTS

(57) Abstract: This invention relates to potent selective agonists of the EP<sub>4</sub> subtype of prostaglandin E<sub>2</sub> receptors, their use or a formulation thereof in the treatment of glaucoma and other conditions which are related to elevated intraocular pressure in the eye of a patient. This invention further relates to the use of the compounds of this invention for mediating the bone modeling and remodeling processes of the osteoblasts and osteoclasts.

WO 2004/037786 A2

BEST AVAILABLE COPY

TITLE OF THE INVENTION  
EP<sub>4</sub> RECEPTOR AGONISTS

CROSS REFERENCE TO RELATED APPLICATIONS

5 This case claims the benefit of provisional application USSN 60/421,402, filed October 25, 2002.

BACKGROUND OF THE INVENTION

) Glaucoma is a degenerative disease of the eye wherein the intraocular pressure is too high to permit normal eye function. As a result, damage may occur to the optic nerve head and result in irreversible loss of visual function. If untreated, glaucoma may eventually lead to blindness. Ocular hypertension, i.e., the condition of elevated intraocular pressure without optic nerve head damage or characteristic glaucomatous visual field defects, is now believed by the majority of ophthalmologists to represent merely the earliest phase in the onset of glaucoma.

5 Many of the drugs formerly used to treat glaucoma proved unsatisfactory. Early methods of treating glaucoma employed pilocarpine and produced undesirable local effects that made this drug, though valuable, unsatisfactory as a first line drug. More recently, clinicians have noted that many  $\beta$ -adrenergic antagonists are effective in reducing intraocular pressure. While many of these agents are effective for this purpose, there exist some patients with whom  
) this treatment is not effective or not sufficiently effective. Many of these agents also have other characteristics, e.g., membrane stabilizing activity, that become more apparent with increased doses and render them unacceptable for chronic ocular use and can also cause cardiovascular effects.

Agents referred to as carbonic anhydrase inhibitors decrease the formation of  
) aqueous humor by inhibiting the enzyme carbonic anhydrase. While such carbonic anhydrase inhibitors are now used to treat elevated intraocular pressure by systemic and topical routes, current therapies using these agents, particularly those using systemic routes are still not without undesirable effects. Topically effective carbonic anhydrase inhibitors are disclosed in U.S. Patent Nos. 4,386,098; 4,416,890; 4,426,388; 4,668,697; 4,863,922; 4,797,413; 5,378,703,  
) 5,240,923 and 5,153,192.

Prostaglandins and prostaglandin derivatives are also known to lower intraocular pressure. There are several prostaglandin types, including the A, B, C, D, E, F, G, I and J- Series (EP 0561073 A1). U.S. Patent 4,883,819 to Bito describes the use and synthesis of PGAs, PGBs and PGCs in reducing intraocular pressure. U.S. Patent 4,824,857 to Goh et al. describes the use  
) and synthesis of PGD<sub>2</sub> and derivatives thereof in lowering intraocular pressure including

derivatives wherein C-10 is replaced with nitrogen. U.S. Patent 5,001,153 to Ueno et al. describes the use and synthesis of 13,14-dihydro-15-keto prostaglandins and prostaglandin derivatives to lower intraocular pressure. U.S. Patent 4,599,353 describes the use of eicosanoids and eicosanoid derivatives including prostaglandins and prostaglandin inhibitors in lowering intraocular pressure. See also WO 00/38667, WO 99/32441, WO 99/02165, WO 00/38663, WO 01/46140, EP 0855389, JP 2000-1472, US Patent No. 6,043,275 and WO 00/38690.

Prostaglandin and prostaglandin derivatives are known to lower intraocular pressure by increasing uveoscleral outflow. This is true for both the F type and A type of prostaglandins. This invention is particularly interested in those compounds that lower IOP via the uveoscleral outflow pathway and other mechanisms by which the E series prostaglandins (PGE<sub>2</sub>) may facilitate IOP reduction. The four recognized subtypes of the EP receptor are believed to modulate the effect of lowering IOP (EP<sub>1</sub>, EP<sub>2</sub>, EP<sub>3</sub> and EP<sub>4</sub>; *J. Lipid Mediators Cell Signaling*, Vol. 14, pages 83-87 (1996)). See also *J. Ocular Pharmacology*, Vol. 4, 1, pages 13-18 (1988); *J. Ocular Pharmacology and Therapeutics*, Vol. 11, 3, pages 447-454 (1995); *J. Lipid Mediators*, Vol. 6, pages 545-553 (1993); US Patent Nos. 5,698,598 and 5,462,968 and *Investigative Ophthalmology and Visual Science*, Vol. 31, 12, pages 2560-2567 (1990). Of particular interest to this invention are compounds, which are agonist of the EP<sub>4</sub> subtype receptor.

A problem with using prostaglandins or derivatives thereof to lower intraocular pressure is that these compounds often induce an initial increase in intraocular pressure, can change the color of eye pigmentation and cause proliferation of some tissues surrounding the eye.

As can be seen, there are several current therapies for treating glaucoma and elevated intraocular pressure, but the efficacy and the side effect profiles of these agents are not ideal. Therefore, there still exist the need for new and effective therapies with little or no side effects.

A variety of disorders in humans and other mammals involve or are associated with abnormal or excessive bone loss. Such disorders include, but are not limited to, osteoporosis, glucocorticoid induced osteoporosis, Paget's disease, abnormally increased bone turnover, periodontal disease, tooth loss, bone fractures, rheumatoid arthritis, periprosthetic osteolysis, osteogenesis imperfecta, metastatic bone disease, hypercalcemia of malignancy, and multiple myeloma. One of the most common of these disorders is osteoporosis, which in its most frequent manifestation occurs in postmenopausal women. Osteoporosis is a systemic skeletal disease characterized by a low bone mass and microarchitectural deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture. Osteoporotic fractures are a major cause of morbidity and mortality in the elderly population. As many as 50%

of women and a third of men will experience an osteoporotic fracture. A large segment of the older population already has low bone density and a high risk of fractures. There is a significant need to both prevent and treat osteoporosis and other conditions associated with bone resorption. Because osteoporosis, as well as other disorders associated with bone loss, are generally chronic conditions, it is believed that appropriate therapy will typically require chronic treatment.

Two different types of cells called osteoblasts and osteoclasts are involved in the bone formation and resorption processes, respectively. See H. Fleisch, *Bisphosphonates In Bone Disease, From The Laboratory To The Patient*, 3rd Edition, Parthenon Publishing (1997), which is incorporated by reference herein in its entirety.

Osteoblasts are cells that are located on the bone surface. These cells secrete an osseous organic matrix, which then calcifies. Substances such as fluoride, parathyroid hormone, and certain cytokines such as prostaglandins are known to provide a stimulatory effect on osteoblast cells. However, an aim of current research is to develop therapeutic agents that will selectively increase or stimulate the bone formation activity of the osteoblasts.

Osteoclasts are usually large multinucleated cells that are situated either on the surface of the cortical or trabecular bone or within the cortical bone. The osteoclasts resorb bone in a closed, sealed-off microenvironment located between the cell and the bone. The recruitment and activity of osteoclasts is known to be influenced by a series of cytokines and hormones. It is well known that bisphosphonates are selective inhibitors of osteoclastic bone resorption, making these compounds important therapeutic agents in the treatment or prevention of a variety of systemic or localized bone disorders caused by or associated with abnormal bone resorption. However, despite the utility of bisphosphonates there remains the desire amongst researchers to develop additional therapeutic agents for inhibiting the bone resorption activity of osteoclasts.

Prostaglandins such as the PGE<sub>2</sub> series are known to stimulate bone formation and increase bone mass in mammals, including man. It is believed that the four different receptor subtypes, designated EP<sub>1</sub>, EP<sub>2</sub>, EP<sub>3</sub>, and EP<sub>4</sub> are involved in mediating the bone modeling and remodeling processes of the osteoblasts and osteoclasts. The major prostaglandin receptor in bone is EP<sub>4</sub>, which is believed to provide its effect by signaling via cyclic AMP.

In present invention it is further found that the formula I agonists of the EP<sub>4</sub> subtype receptor are useful for stimulating bone formation.


WO 02/24647, WO 02/42268, EP 1132086, EP 855389, EP 1114816, WO 01/46140 and WO 01/72268 disclose EP<sub>4</sub> agonists.

## SUMMARY OF THE INVENTION

This invention relates to potent selective agonists of the EP4 subtype of prostaglandin E2 receptors, formulations thereof, and their use in the treatment of glaucoma and other conditions that are related to elevated intraocular pressure in the eye of a patient. The invention also relates to the use of the compounds to provide a neuroprotective effect to the eye of mammalian species, particularly humans. The invention also relates to the use of the compounds for mediating the bone modeling and remodeling processes of the osteoblasts and osteoclasts.

More particularly, this invention relates to novel EP4 agonist having the structural

formula I:



FORMULA I

or a pharmaceutically acceptable salt thereof, wherein,

Y¹ is

1) CH<sub>2</sub>CH<sub>2</sub>,

2) CHCH, or



Y is C(O) or CH(OH);

A is (CH<sub>2</sub>)<sub>n</sub>;

n is 1, 2, 3, or 4;

W a bond, unsubstituted C 1-6 alkylene, or C 1-6 alkylene substituted with 1, 2, 3, or 4 halogen atoms;

Z is

1) O,

2) S,




4) HC=CH,


5)  $C \equiv C$ , or

6) a bond;

Q is a disubstituted aryl or heteroaryl ring, wherein one ring atom of the ring is attached to the moiety



and another ring atom is attached to the moiety



$R^1$  is

$COR^5$ ,

$OH$ ,

$CN$ ,

$(CH_2)_{1-3} CO_2R^6$ ,

$C(O)NHSO_2R^8$ ,

$SO_2R^7$ ,

$(CH_2)_{0-4} SO_3R^6$ ,

$CF_2SO_2NH_2$ ,

$SO_2NH_2$ ,

$SO_2NHCOR^8$ ,

$PO(OR^7)_2$ ,

$C_{1-4}$  alkoxy,

hydroxymethylketone, or

$(CH_2)_{0-4} R^k$ , wherein  $R^k$  is unsubstituted or substituted with 1 to 3 groups of  $R^a$ ;

$R^2$  is

1)  $C_{1-6}$  alkyl,

2)  $(CH_2)_{0-8} C_{6-10}$  aryl,

3)  $(CH_2)_{0-8} R^m$ ,

4)  $(CH_2)_{0-8} C_{3-8}$  cycloalkyl,

5)  $O-C_{1-10}$  alkyl,

6)  $O-C_{6-10}$  aryl,

7)  $O-R^m$ ,

8)  $O-C_{3-10}$  cycloalkyl

wherein aryl,  $R^m$ , and cycloalkyl are unsubstituted or substituted with 1-3 groups of  $R^b$ ;

$R^3$  and  $R^4$  are independently selected from the group consisting of

- 1) halogen, and
- 2)  $C_{1-6}$  alkyl, or

5  $R^3$  and  $R^4$ , together with the carbon atom to which they are attached, form a  $C_{3-7}$  cycloalkyl ring;

$R^5$  is

- 1) hydrogen,
- 2) OH,
- 10 3)  $CH_2OH$ ,
- 4)  $C_{1-6}$  alkoxy,
- 5)  $NHPO_2R^6$ ,
- 6)  $NHR^9$ ,
- 7)  $NHSO_2R^8$ , or
- 15 8)  $NR^6R^7$ ;

$R^6$  and  $R^7$  are independently selected from the group consisting of hydrogen,  $C_{1-6}$  alkyl, and  $C_{3-8}$  cycloalkyl;

$R^8$  is selected from the group consisting of hydrogen,  $C_{6-10}$  aryl,  $R^n$ , and  $C_{1-4}$  alkyl;

$R^9$  is  $C(O)R^{10}$  or  $SO_2R^{10}$ ;

20  $R^{10}$  is hydrogen,  $C_{6-10}$  aryl, or  $C_{1-4}$  alkyl;

$R^a$  and  $R^b$  are independently selected from the group consisting of

- 1)  $C_{1-6}$  alkoxy,
- 2)  $C_{1-6}$  alkyl, unsubstituted or substituted with
  - a)  $C_{1-6}$  alkoxy,
  - 25 b)  $C_{1-6}$  alkylthio,
  - c) CN,
  - d) OH, or
  - e)  $CF_3$ ,
- 3)  $CF_3$ ,
- 30 4) nitro,
- 5) amino,
- 6) cyano,
- 7)  $C_{1-6}$  alkylamino,
- 8) halogen
- 35 9) ORC,

10)  $\text{OCH}_2\text{R}^c$ , and

11)  $\text{CH}_2\text{OR}^c$ ;

$\text{R}^c$  is

1)  $\text{C}_6$ -10aryl,

2)  $\text{R}^s$ , or

3)  $\text{C}_3$ -8cycloalkyl; and

$\text{R}^k$ ,  $\text{R}^m$ ,  $\text{R}^n$  and  $\text{R}^s$  are independently selected from the group consisting of

1) a stable monocyclic heteroaryl ring having 5, 6 or 7 ring atoms, or a stable bicyclic heteroaryl ring having 8, 9, 10, or 11 ring atoms, wherein the monocyclic ring has 1, 2, 3, or 4 heteroatoms, independently selected from the group consisting of O, S or N, and wherein the bicyclic ring has 1, 2, 3, or 4 heteroatoms, independently selected from the group consisting of O, S or N, and

2) a stable monocyclic or bicyclic heterocycloalkyl ring system a stable, saturated monocyclic or bicyclic ring system having 3 to 10 ring atoms, wherein 1, 2, 3, or 4 ring atoms are heteroatoms selected from O, S and N.

The compounds of the present invention may have chiral centers and occur as racemates, racemic mixtures and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention. The compounds of the present invention may also have polymorphic crystalline forms, with all polymorphic crystalline forms being included in the present invention. The compounds of the invention also include tautomeric forms, with all tautomeric forms being included in the present invention.

The invention also includes prodrug forms of the above-described compounds. Prodrugs, such as ester derivatives of active drug, are compound derivatives which, when absorbed into the bloodstream of a warm-blooded animal, cleave in such a manner as to release the drug form and permit the drug to afford improved therapeutic efficacy. The prodrugs may be administered in low amounts relative to the amounts of antagonist that would ordinarily be administered. The prodrugs may be administered orally. The prodrugs retain structural integrity while passing through the gastrointestinal system, and are effectively delivered to cells. They are subjected to metabolic reactions to form the active acid which then interacts with the platelet receptor site.

This and other aspects of the invention will be realized upon inspection of the invention as a whole.




## DETAILED DESCRIPTION OF THE INVENTION

In a class of compounds of the invention, and pharmaceutically acceptable salts thereof, Y<sup>1</sup> is CHCH and Y is CH(OH).


In a subclass of this class, A is (CH<sub>2</sub>)<sub>1-3</sub> and W is a bond or (CH<sub>2</sub>)<sub>1-3</sub>.

- 5 In a group of this subclass, 1) R<sup>1</sup> is COOH or tetrazole, 2) R<sup>2</sup> is phenyl, and 3) R<sup>3</sup> and R<sup>4</sup> are halogen, or R<sup>3</sup> and R<sup>4</sup> together with the carbon to which they are attached, form a cyclopropyl ring.

In a subgroup of this group, Q is selected from the group consisting of








In a family of this subgroup, Q is selected from the group consisting of



- 15 In the above family of structures, "(Z)" and "(W)" indicate the atoms in Q to which variables Z and W defined above are attached.

Examples of the subgroup include





- 5 (1) 5-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)thiophene-2-carboxylic acid
- (2) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[5-(1*H*-tetrazol-5-yl)thien-2-yl]propyl}pyrrolidin-2-one
- 10 (3) 5-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)-1,3-thiazole-2-carboxylic acid

- (4) 2-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)-1,3-thiazole-5-carboxylic acid
- 5 (5) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[5-(1*H*-tetraazol-5-yl)-1,3-thiazol-2-yl]propyl}pyrrolidin-2-one
- (6) 2-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)-1,3-thiazole-4-carboxylic acid
- 10 (7) [5-(2-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)ethyl)thien-2-yl]acetic acid
- (8) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{2-[5-(1*H*-tetraazol-5-yl)methyl]thien-2-yl}ethyl}pyrrolidin-2-one
- 15 (9) 2-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)-1,3-oxazole-5-carboxylic acid
- (10) 5-(3-((2*R*)-2-[(1*E*)-3-hydroxy-3-(1-phenylcyclopropyl)prop-1-enyl]-5-oxopyrrolidin-1-yl)propyl)thiophene-2-carboxylic acid
- 20 (11) (5*R*)-5-[(1*E*)-3-hydroxy-3-(1-phenylcyclopropyl)prop-1-enyl]-1-{3-[5-(1*H*-tetraazol-5-yl)thien-2-yl]propyl}pyrrolidin-2-one
- 25 (12) 5-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)-1,3,4-thiadiazole-2-carboxylic acid
- (13) 4-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)benzoic acid
- 30 (14) 3-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)benzoic acid

- (15) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[3-(1H-tetraazol-5-yl)phenyl]propyl}pyrrolidin-2-one
- 5 (16) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[4-(1H-tetraazol-5-yl)phenyl]propyl}pyrrolidin-2-one
- (17) 3-[5-((2R)-2-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)methyl]thien-2-yl]propanoic acid
- 10 (18) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-({5-[2-(1H-tetraazol-5-yl)ethyl]thien-2-yl)methyl}pyrrolidin-2-one
- (19) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-({4-[3-(1H-tetraazol-5-yl)propyl]thien-3-yl)methyl}pyrrolidin-2-one
- 15 (20) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-({4-[2-(1H-tetraazol-5-yl)ethyl]thien-2-yl)methyl}pyrrolidin-2-one
- (21) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-({4-[2-(1H-tetraazol-5-yl)ethyl]-1,3-thiazol-2-yl)methyl}pyrrolidin-2-one
- 20 (22) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[2-(1H-tetraazol-5-yl)ethyl]benzyl}pyrrolidin-2-one

25 The invention is described herein in detail using the terms defined below unless otherwise specified.

30 The term "therapeutically effective amount", as used herein, means that amount of the EP<sub>4</sub> receptor subtype agonist of formula I, or other actives of the present invention, that will elicit the desired therapeutic effect or response or provide the desired benefit when administered in accordance with the desired treatment regimen. A preferred therapeutically effective amount relating to the treatment of abnormal bone resorption is a bone formation, stimulating amount. Likewise, a preferred therapeutically effective amount relating to the treatment of ocular hypertension or glaucoma is an amount effective for reducing intraocular pressure and/or treating ocular hypertension and/or glaucoma.

The term "pharmaceutically acceptable" as used herein, means generally suitable for administration to a mammal, including humans, from a toxicity or safety standpoint.

The term "prodrug" refers to compounds which are drug precursors which, following administration and absorption, release the claimed drug in vivo via some metabolic process. A non-limiting example of a prodrug of the compounds of this invention would be an acid of the pyrrolidinone group, where the acid functionality has a structure that makes it easily hydrolyzed after administration to a patient. Exemplary prodrugs include acetic acid derivatives that are non-narcotic, analgesics/non-steroidal, anti-inflammatory drugs having a free  $\text{CH}_2\text{COOH}$  group (which can optionally be in the form of a pharmaceutically acceptable salt, e.g.  $-\text{CH}_2\text{COO}-\text{Na}^+$ ), typically attached to a ring system, preferably to an aromatic or heteroaromatic ring system.

The term "alkyl", unless otherwise specified, refers to a monovalent alkane (hydrocarbon) derived radical containing from 1 to 10 carbon atoms unless otherwise defined. It may be straight, branched or cyclic. Preferred alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, t-butyl, cyclopentyl and cyclohexyl. When the alkyl group is said to be substituted with an alkyl group, this is used interchangeably with "branched alkyl group". Corresponding divalent groups are referred to as "alkylene" groups, e.g. methylene, ethylene, etc.

Variables which include alkenylenes such as ethenylene (e.g.  $-\text{CH}=\text{CH}-$ ), unless otherwise specified, are represented by "CHCH". The term "alkoxy" refers to  $\text{C}_1\text{-C}_6$  alkyl-O-, with the alkyl group optionally substituted as described herein. Examples of alkoxy groups are methoxy, ethoxy, propoxy, butoxy and isomeric groups thereof.

The terms "halogen" or "halo" refer to chlorine, fluorine, iodine or bromine.

The term "aryl" refers to aromatic rings e.g., phenyl, substituted phenyl and the like, as well as rings which are fused, e.g., naphthyl, phenanthrenyl and the like. An aryl group thus contains at least one ring having at least 6 atoms, with up to five such rings being present, containing up to 22 atoms therein, with alternating (resonating) double bonds between adjacent carbon atoms or suitable heteroatoms. The preferred aryl groups are phenyl, naphthyl and phenanthrenyl. Unless otherwise specified, the aryl ring can be unsubstituted or substituted with one or more of  $-\text{CF}_3$ ,  $-\text{CN}$ ,  $\text{C}_{1-4}$  alkyl, hydroxy,  $\text{C}_{1-4}$  alkoxy, halogen, e.g. F, Cl, Br, or I,  $-\text{NO}_2$ ,  $-\text{NR}^{\text{d}}\text{R}^{\text{f}}$ ,  $-\text{SO}_2\text{R}^{\text{d}}$ ,  $\text{SO}_2\text{NR}^{\text{d}}\text{R}^{\text{f}}$ ,  $-\text{CONR}^{\text{d}}\text{R}^{\text{f}}$ , or  $\text{COR}^{\text{d}}$ , wherein  $\text{R}^{\text{d}}$  and  $\text{R}^{\text{f}}$  are independently selected hydrogen and  $\text{C}_{1-4}$  alkyl. Preferred substituted aryls include phenyl and naphthyl.

The term "heterocycloalkyl", unless otherwise specified, refers to a stable, saturated monocyclic or bicyclic ring system having 3 to 10 ring atoms, wherein 2 to 6 ring atoms are carbon atoms, and 1 to 4 ring atoms are heteroatoms selected from O, S and N. Unless

otherwise specified, the heterocycloalkyl ring can be unsubstituted or substituted with one or more of C<sub>1-4</sub> alkyl, hydroxy, C<sub>1-4</sub> alkoxy, amino, and halogen, e.g. F, Cl, Br, or I.

The term "cycloalkyl", unless otherwise specified, refers to a cyclic alkyl group (nonaromatic) having the specified number of carbon atoms, e.g., C<sub>3-7</sub> cycloalkyl has 3, 4, 5, 6,

5 or 7 carbon atoms. Unless otherwise specified, the cycloalkyl ring can be unsubstituted or substituted with one or more of C<sub>1-4</sub> alkyl, hydroxy, C<sub>1-4</sub> alkoxy, amino, and halogen, e.g. F, Cl, Br, or I. Examples include cyclopropyl, cyclobutyl, and cyclopentyl.

The term "heteroatom" means O, S or N, selected on an independent basis.

10 The term "heteroaryl", unless otherwise specified, refers to an unsaturated monocyclic aromatic hydrocarbon group having 5, 6 or 7 ring atoms, or an unsaturated bicyclic aromatic group having 8, 9, 10, or 11 ring atoms, containing 1, 2, 3, or 4 heteroatoms, independently selected from the group consisting of O, S or N, in which a carbon or nitrogen atom is the point of attachment. Examples of this type are pyrrole, pyridine, oxazole, thiazole, tetrazole, and oxazine. Unless otherwise specified, the heteroaryl ring can be unsubstituted or  
15 substituted with one or more of C<sub>1-4</sub> alkyl, hydroxy, C<sub>1-4</sub> alkoxy, amino, and halogen, e.g. F, Cl, Br, or I. For purposes of this invention the tetrazole includes all tautomeric forms. Additional nitrogen atoms may be present together with the first nitrogen and oxygen or sulfur, giving, e.g., thiadiazole.


Bicyclic heteroaryl rings include bicyclic ring systems in which either or both  
20 rings contain heteroatoms. Included within, but not limiting this term, are systems in which one ring contains 1, 2, 3, or 4 heteroatoms and the other ring is a benzene ring.

Bicyclic heterocycloalkyl rings include bicyclic ring systems in which either or both rings contain heteroatoms. Included within, but not limiting this term, are systems in which one ring contains 1, 2, 3, or 4 heteroatoms and the other ring contains zero heteroatoms.


25 The heterocycloalkyl or heteroaryl ring may be attached at any heteroatom or carbon atom that results in the creation of a stable structure. Examples of such rings include, but are not limited to, azepinyl, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothieryl, benzoxazolyl, chromanyl, cinnolinyl, dihydrobenzofuryl, dihydrobenzothieryl, dihydrobenzothiopyranyl,  
30 dihydrobenzothiopyranyl sulfone, 1,3-dioxolanyl, furyl, imidazolidinyl, imidazolinyl, imidazolyl, indolinyl, indolyl, isochromanyl, isoindolinyl, isoquinolinyl, isothiazolidinyl, isothiazolyl, isothiazolidinyl, morpholinyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, 2-oxopiperazinyl, 2-oxopiperdinyl, 2-oxopyrrolidinyl, piperidyl, piperazinyl, pyridyl, pyrazinyl, pyrazolidinyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrrolidinyl, pyrrolyl, quinazolinyl, quinolinyl,

quinoxaliny, tetrahydrofuryl, tetrahydroisoquinoliny, tetrahydroquinoliny, thiamorpholiny, thiamorpholiny sulfoxide, thiazolyl, thiazoliny, thienofuryl, thienothieryl, thienyl, and triazolyl.

The term "a disubstituted aryl or heteroaryl ring" includes aryl and heteroaryl rings in which two ring carbon atoms have substituents attached and do not have hydrogen atoms attached, e.g. 2,5-substituted thiophene, furan, and thiazole, and 1,2-, 1,3- and 1,4-substituted benzene. Such disubstituted rings include, but are not limited to, those structurally depicted as




In a preferred embodiment, the disubstituted aryl ring is




In another preferred embodiment, the disubstituted heteroaryl ring is



In another preferred embodiment, the disubstituted heteroaryl ring is




In another preferred embodiment, the disubstituted heteroaryl ring is




In another preferred embodiment, the disubstituted heteroaryl ring is






In another preferred embodiment, the disubstituted heteroaryl ring is




In another preferred embodiment, the disubstituted heteroaryl ring is




5

In another preferred embodiment, the disubstituted heteroaryl ring is




In another preferred embodiment, the disubstituted heteroaryl ring is




10


In another preferred embodiment, the disubstituted heteroaryl ring is



In another preferred embodiment, the disubstituted heteroaryl ring is




In another preferred embodiment, the disubstituted heteroaryl ring is




15

In another preferred embodiment, the disubstituted heteroaryl ring is



When Q is defined to include substituted heteroaryl rings shown as



"(Z)" and "(W)" represent variables "Z" and "W", and are presented to clearly identify the atom in Q to which these variables are attached.

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced.

The term "agonist" as used herein means EP4 subtype compounds of formula I interact with the EP4 receptor to produce maximal, super maximal or submaximal effects compared to the natural agonist, PGE2. See Goodman and Gilman, *The Pharmacological Basis of Therapeutics*, 9<sup>th</sup> edition, 1996, chapter 2.

Another embodiment of this invention is directed to a composition containing an EP4 agonist of Formula I and optionally a pharmaceutically acceptable carrier.

Yet another embodiment of this invention is directed to a method for decreasing elevated intraocular pressure or treating glaucoma by administration, preferably topical or intracameral administration, of a composition containing an EP4 agonist of Formula I and optionally a pharmaceutically acceptable carrier. Use of the compounds of formula I for the manufacture of a medicament for treating elevated intraocular pressure or glaucoma or a combination thereof is also included in this invention.

This invention is further concerned with a process for making a pharmaceutical composition comprising a compound of formula I.

This invention is further concerned with a process for making a pharmaceutical composition comprising a compound of formula I, and a pharmaceutically acceptable carrier.

The claimed compounds bind strongly and act on PGE<sub>2</sub> receptor, particularly on the EP4 subtype receptor and therefore are useful for preventing and/or treating glaucoma and ocular hypertension.

Dry eye is a common ocular surface disease afflicting millions of people. Although it appears that dry eye may result from a number of unrelated pathogenic causes, the common end result is the breakdown of the tear film, which results in dehydration of the exposed outer surface of the eye. (Lemp, Report of the Nation Eye Institute/Industry Workshop on

Clinical Trials in Dry Eyes, The *CLAO Journal*, 21(4):221-231 (1995)). One cause for dry eye is the decreased mucin production by the conjunctival cells and/or corneal epithelial cells of mucin, which protects and lubricates the ocular surface (Gipson and Inatomi, Mucin genes expressed by ocular surface epithelium. *Progress in Retinal and Eye Research*, 16:81-98 (1997)). Functional  
5 EP4 receptors have been found in human conjunctival epithelial cells (see US Patent 6,344,477, incorporated by reference in its entirety) and it is appreciated that both human corneal epithelial cells (*Progress in Retinal and Eye Research*, 16:81-98(1997)) and conjunctival cells (Dartt et al. Localization of nerves adjacent to goblet cells in rat conjunctiva. *Current Eye Research*, 14:993-1000 (1995)) are capable of secreting mucins. Thus, the compounds of formula I are useful for  
10 treating dry eye.

Macular edema is swelling within the retina within the critically important central visual zone at the posterior pole of the eye. An accumulation of fluid within the retina tends to detach the neural elements from one another and from their local blood supply, creating a dormancy of visual function in the area. It is believed that EP4 agonist which lower IOP are  
15 useful for treating diseases of the macular such as macular edema or macular degeneration. Thus, another aspect of this invention is a method for treating macular edema or macular degeneration.

Glaucoma is characterized by progressive atrophy of the optic nerve and is frequently associated with elevated intraocular pressure (IOP). It is possible  
20 to treat glaucoma, however, without necessarily affecting IOP by using drugs that impart a neuroprotective effect. See *Arch. Ophthalmol.* Vol. 112, Jan 1994, pp. 37-44; *Investigative Ophthalmol. & Visual Science*, 32, 5, April 1991, pp. 1593-99. It is believed that EP4 agonist which lower IOP are useful for providing a neuroprotective effect. They are also believed to be effective for increasing retinal and optic nerve head blood velocity and  
25 increasing retinal and optic nerve oxygen by lowering IOP, which when coupled together benefits optic nerve health. As a result, this invention further relates to a method for increasing retinal and optic nerve head blood velocity, or increasing retinal and optic nerve oxygen tension or providing a neuroprotective effect or a combination thereof by using an EP4 agonist of formula I.

The compounds produced in the present invention are readily combined with  
30 suitable and known pharmaceutically acceptable excipients to produce compositions which may be administered to mammals, including humans, to achieve effective IOP lowering. Thus, this invention is also concerned with a method of treating ocular hypertension or glaucoma by administering to a patient in need thereof one of the compounds of formula I alone or in combination with a  $\beta$ -adrenergic blocking agent such as timolol, betaxolol, levobetaxolol,  
35 carteolol, levobunolol, a parasympathomimetic agent such as pilocarpine, a sympathomimetic

agents such as epinephrine, iopidine, brimonidine, clonidine, para-aminoclonidine, a carbonic anhydrase inhibitor such as dorzolamide, acetazolamide, metazolamide or brinzolamide; a prostaglandin such as latanoprost, travaprost, unoprostone, rescula, S1033 (compounds set forth in US Patent Nos. 5,889,052; 5,296,504; 5,422,368; and 5,151,444); a hypotensive lipid such as lumigan and the compounds set forth in US Patent No. 5,352,708; a neuroprotectant disclosed in US Patent No. 4,690,931, particularly eliprodil and R-eliprodil as set forth in WO 94/13275, including memantine; or an agonist of 5-HT<sub>2</sub> receptors as set forth in PCT/US00/31247, particularly 1-(2-aminopropyl)-3-methyl-1H-imidazol-6-ol fumarate and 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-ethylamine.

This invention is also concerned with a method for increasing retinal and optic nerve head blood velocity, or increasing retinal and optic nerve oxygen tension or providing a neuroprotective effect or a combination thereof by administering to a patient in need thereof one of the compounds of formula I alone or in combination with a  $\beta$ -adrenergic blocking agent such as timolol, betaxolol, levobetaxolol, carteolol, levobunolol, a parasympathomimetic agent such as pilocarpine, a sympathomimetic agents such as epinephrine, iopidine, brimonidine, clonidine, para-aminoclonidine, a carbonic anhydrase inhibitor such as dorzolamide, acetazolamide, metazolamide or brinzolamide; a prostaglandin such as latanoprost, travaprost, unoprostone, rescula, S1033 (compounds set forth in US Patent Nos. 5,889,052; 5,296,504; 5,422,368; and 5,151,444); a hypotensive lipid such as lumigan and the compounds set forth in US Patent No. 5,352,708; a neuroprotectant disclosed in US Patent No. 4,690,931, particularly eliprodil and R-eliprodil as set forth in WO 94/13275, including memantine; or an agonist of 5-HT<sub>2</sub> receptors as set forth in PCT/US00/31247, particularly 1-(2-aminopropyl)-3-methyl-1H-imidazol-6-ol fumarate and 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-ethylamine. Use of the compounds of formula I for the manufacture of a medicament for increasing retinal and optic nerve head blood velocity, or increasing retinal and optic nerve oxygen tension or providing a neuroprotective effect or a combination thereof is also included in this invention.

This invention is further concerned with a method for treating macular edema or macular degeneration by administering to a patient in need thereof one of the compounds of formula I alone or in combination with a  $\beta$ -adrenergic blocking agent such as timolol, betaxolol, levobetaxolol, carteolol, levobunolol, a parasympathomimetic agent such as pilocarpine, a sympathomimetic agents such as epinephrine, iopidine, brimonidine, clonidine, para-aminoclonidine, a carbonic anhydrase inhibitor such as dorzolamide, acetazolamide, metazolamide or brinzolamide; a prostaglandin such as latanoprost, travaprost, unoprostone, rescula, S1033 (compounds set forth in US Patent Nos. 5,889,052; 5,296,504; 5,422,368; and 5,151,444); a hypotensive lipid such as lumigan and the compounds set forth in US Patent No.

5,352,708; a neuroprotectant disclosed in US Patent No. 4,690,931, particularly eliprodil and R-eliprodil as set forth in WO 94/13275, including memantine; a Maxi-K channel blocker such as Penitrem A, paspalicine, charybdotoxin, iberiotoxin or as disclosed in USSN 60/389,205, filed June 17, 2002 (Attorney Docket 21121PV), 60/389,222, filed June 17, 2002 (Attorney docket 21092PV), 60/458,981, filed March 27, 2003 (Attorney docket 21101PV4), 60/424790, filed November 8, 2002 (Attorney docket 21260PV), 60/424808, filed November 8, 2002 (Attorney docket 21281PV), 09/765716, filed January 17, 2001, 09/764738, filed January 17, 2001 and PCT publications WO 02/077168 and WO 02/02060863, all incorporated by reference in their entirety herein; or an agonist of 5-HT<sub>2</sub> receptors as set forth in PCT/US00/31247, particularly 1-(2-aminopropyl)-3-methyl-1H-imidazol-6-ol fumarate and 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-ethylamine. Use of the compounds of formula I for the manufacture of a medicament for macular edema or macular degeneration is also included in this invention.

Compounds of the invention may also be used to treat neuropathic pain.

Neuropathic pain syndromes can develop following neuronal injury and the resulting pain may persist for months or years, even after the original injury has healed. Neuronal injury may occur in the peripheral nerves, dorsal roots, spinal cord or certain regions in the brain. Neuropathic pain syndromes are traditionally classified according to the disease or event that precipitate them. Neuropathic pain syndromes include: diabetic neuropathy; sciatica; non-specific lower back pain; multiple sclerosis pain; fibromyalgia; HIV-related neuropathy, post-herpetic neuralgia; trigeminal neuralgia; and pain resulting from physical trauma, amputation, cancer, toxins or chronic inflammatory conditions. These conditions are difficult to treat and although several drugs are known to have limited efficacy, complete pain control is rarely achieved. The symptoms of neuropathic pain are incredibly heterogeneous and are often described as spontaneous shooting and lancinating pain, or ongoing, burning pain. In addition, there is pain associated with normally non-painful sensations such as "pins and needles" (paraesthesias and dysesthesias), increased sensitivity to noxious stimuli (thermal, cold, mechanical hyperalgesia), continuing pain sensation after removal of the stimulation (hyperpathia) or an absence of or deficit in selective sensory pathways (hypoalgesia).

Compounds of the invention may also be used to treat acute renal failure, chronic renal failure, colon cancer, colitis, and HIV latency.

The EP<sub>4</sub> agonist used in the instant invention can be administered in a therapeutically effective amount intravenously, subcutaneously, topically, transdermally, parenterally or any other method known to those skilled in the art. Ophthalmic pharmaceutical compositions are preferably adapted for topical administration to the eye in the form of solutions, suspensions, ointments, creams or

as a solid insert. Ophthalmic formulations of this compound may contain from 0.001 to 5% and especially 0.001 to 0.1% of medicament. Higher dosages as, for example, up to about 10% or lower dosages can be employed provided the dose is effective in reducing intraocular pressure, treating glaucoma, increasing blood flow velocity or oxygen tension. For a single dose, from  
5 between 0.001 to 5.0 mg, preferably 0.005 to 2.0 mg, and especially 0.005 to 1.0 mg of the compound can be applied to the human eye.

The pharmaceutical preparation which contains the compound may be conveniently admixed with a non-toxic pharmaceutical organic carrier, or with a non-toxic pharmaceutical inorganic carrier. Typical of pharmaceutically acceptable carriers are, for  
10 example, water, mixtures of water and water-miscible solvents such as lower alkanols or aralkanols, vegetable oils, peanut oil, polyalkylene glycols, petroleum based jelly, ethyl cellulose, ethyl oleate, carboxymethyl-cellulose, polyvinylpyrrolidone, isopropyl myristate and other conventionally employed acceptable carriers. The pharmaceutical preparation may also contain non-toxic auxiliary substances such as emulsifying, preserving,  
15 wetting agents, bodying agents and the like, as for example, polyethylene glycols 200, 300, 400 and 600, carbowaxes 1,000, 1,500, 4,000, 6,000 and 10,000, antibacterial components such as quaternary ammonium compounds, phenylmercuric salts known to have cold sterilizing properties and which are non-injurious in use, thimerosal, methyl and propyl paraben, benzyl alcohol, phenyl ethanol, buffering ingredients such as sodium borate, sodium acetates, gluconate  
20 buffers, and other conventional ingredients such as sorbitan monolaurate, triethanolamine, oleate, polyoxyethylene sorbitan monopalmitate, dioctyl sodium sulfosuccinate, monothioglycerol, thiosorbitol, ethylenediamine tetracetic acid, and the like. Additionally, suitable ophthalmic vehicles can be used as carrier media for the present purpose including conventional phosphate buffer vehicle systems, isotonic boric acid vehicles, isotonic sodium chloride vehicles, isotonic sodium borate vehicles and the like. The pharmaceutical preparation may also be in the form of a microparticle formulation. The pharmaceutical preparation may also be in the form of a solid insert. For example, one may use a solid water soluble polymer as the carrier for the medicament. The polymer used to form the insert may be any water soluble non-toxic polymer, for example, cellulose derivatives such as methylcellulose, sodium carboxymethyl cellulose,  
25 (hydroxyloweralkyl cellulose), hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose; acrylates such as polyacrylic acid salts, ethylacrylates, polyacrylamides; natural products such as gelatin, alginates, pectins, tragacanth, karaya, chondrus, agar, acacia; the starch derivatives such as starch acetate, hydroxymethyl starch ethers, hydroxypropyl starch, as well as other synthetic derivatives such as polyvinyl alcohol, polyvinyl

pyrrolidone, polyvinyl methyl ether, polyethylene oxide, neutralized carbopol and xanthan gum, gellan gum, and mixtures of said polymer.

Suitable subjects for the administration of the formulation of the present invention include primates, man and other animals, particularly man and domesticated animals such as cats, rabbits and dogs.

The pharmaceutical preparation may contain non-toxic auxiliary substances such as antibacterial components which are non-injurious in use, for example, thimerosal, benzalkonium chloride, methyl and propyl paraben, benzylidodecylmethyl ammonium bromide, benzyl alcohol, or phenylethanol; buffering ingredients such as sodium chloride, sodium borate, sodium acetate, sodium citrate, or gluconate buffers; and other conventional ingredients such as sorbitan monolaurate, triethanolamine, polyoxyethylene sorbitan monopalmitate, ethylenediamine tetraacetic acid, and the like.

The ophthalmic solution or suspension may be administered as often as necessary to maintain an acceptable IOP level in the eye. It is contemplated that administration to the mammalian eye will be from once up to three times daily.

For topical ocular administration the novel formulations of this invention may take the form of solutions, gels, ointments, suspensions or solid inserts, formulated so that a unit dosage comprises a therapeutically effective amount of the active component or some multiple thereof in the case of a combination therapy.

The compounds of the instant invention are also useful for mediating the bone modeling and remodeling processes of the osteoblasts and osteoclasts. See PCT US99/23757 filed October 12, 1999 and incorporated herein by reference in its entirety. The major prostaglandin receptor in bone is EP<sub>4</sub>, which is believed to provide its effect by signaling via cyclic AMP. See Ikeda T, Miyaura C, Ichikawa A, Narumiya S, Yoshiki S and Suda T 1995, *In situ localization of three subtypes (EP<sub>1</sub>, EP<sub>3</sub> and EP<sub>4</sub>) of prostaglandin E receptors in embryonic and newborn mice.*, *J Bone Miner Res* 10 (sup 1):S172, which is incorporated by reference herein in its entirety. Use of the compounds of formula I for the manufacture of a medicament for mediating the bone modeling and remodeling processes are also included in this invention

Thus, another object of the present invention is to provide methods for stimulating bone formation, i.e. osteogenesis, in a mammal comprising administering to a mammal in need thereof a therapeutically effective amount of an EP<sub>4</sub> receptor subtype agonist of formula I.

Still another object of the present invention to provide methods for stimulating bone formation in a mammal in need thereof comprising administering to said mammal a therapeutically effective amount of an EP<sub>4</sub> receptor subtype agonist of formula I and a

bisphosphonate active. Use of the compounds of formula I for the manufacture of a medicament for stimulating bone formation is also included in this invention.

Yet another object of the present invention to provide pharmaceutical compositions comprising a therapeutically effective amount of an EP<sub>4</sub> receptor subtype agonist of formula I and a bisphosphonate active.

It is another object of the present invention to provide methods for treating or reducing the risk of contracting a disease state or condition related to abnormal bone resorption in a mammal in need of such treatment or prevention, comprising administering to said mammal a therapeutically effective amount of an EP<sub>4</sub> receptor subtype agonist of formula I. Use of the compounds of formula I for the manufacture of a medicament for treating or reducing the risk of contracting a disease state or condition related to abnormal bone resorption is also included in this invention.

The disease states or conditions related to abnormal bone resorption include, but are not limited to, osteoporosis, glucocorticoid induced osteoporosis, Paget's disease, abnormally increased bone turnover, periodontal disease, tooth loss, bone fractures, rheumatoid arthritis, periprosthetic osteolysis, osteogenesis imperfecta, metastatic bone disease, hypercalcemia of malignancy, and multiple myeloma.

Within the method comprising administering a therapeutically effective amount of an EP<sub>4</sub> receptor subtype agonist of formula I and a bisphosphonate active, both concurrent and sequential administration of the EP<sub>4</sub> receptor subtype agonist of formula I and the bisphosphonate active are deemed within the scope of the present invention. Generally, the formulations are prepared containing 5 or 10 mg of a bisphosphonate active, on a bisphosphonic acid active basis. With sequential administration, the agonist and the bisphosphonate can be administered in either order. In a subclass of sequential administration the agonist and bisphosphonate are typically administered within the same 24 hour period. In yet a further subclass, the agonist and bisphosphonate are typically administered within about 4 hours of each other.

Nonlimiting examples of bisphosphonate actives useful herein include the following:

Alendronic acid, 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;

Alendronate (also known as alendronate sodium or alendronate monosodium trihydrate), 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium trihydrate;

Alendronic acid and alendronate are described in U.S. Patents 4,922,007, to Kieczkowski et al., issued May 1, 1990; 5,019,651, to Kieczkowski et al., issued May 28,



1991; 5,510,517, to Dauer et al., issued April 23, 1996; 5,648,491, to Dauer et al., issued July 15, 1997, all of which are incorporated by reference herein in their entirety;

Cycloheptylaminomethylene-1,1-bisphosphonic acid, YM 175, Yamanouchi (cimadronate), as described in U.S. Patent 4,970,335, to Isomura et al., issued November 13, 1990, which is incorporated by reference herein in its entirety;

1,1-dichloromethylene-1,1-diphosphonic acid (clodronic acid), and the disodium salt (clodronate, Procter and Gamble), are described in Belgium Patent 672,205 (1966) and *J. Org. Chem* 32, 4111 (1967), both of which are incorporated by reference herein in their entirety;

1-hydroxy-3-(1-pyrrolidiny)-propylidene-1,1-bisphosphonic acid (EB-1053);  
1-hydroxyethane-1,1-diphosphonic acid (etidronic acid);  
1-hydroxy-3-(N-methyl-N-pentylamino)propylidene-1,1-bisphosphonic acid, also known as BM-210955, Boehringer-Mannheim (ibandronate), is described in U.S. Patent No. 4,927,814, issued May 22, 1990, which is incorporated by reference herein in its entirety;  
6-amino-1-hydroxyhexylidene-1,1-bisphosphonic acid (neridronate);  
3-(dimethylamino)-1-hydroxypropylidene-1,1-bisphosphonic acid (olpadronate);

3-amino-1-hydroxypropylidene-1,1-bisphosphonic acid (pamidronate);  
[2-(2-pyridinyl)ethylidene]-1,1-bisphosphonic acid (piridronate) is described in U.S. Patent No. 4,761,406, which is incorporated by reference in its entirety;  
1-hydroxy-2-(3-pyridinyl)-ethylidene-1,1-bisphosphonic acid (risedronate);  
(4-chlorophenyl)thiomethane-1,1-disphosphonic acid (tiludronate) as described in U.S. Patent 4,876,248, to Breliere et al., October 24, 1989, which is incorporated by reference herein in its entirety; and

1-hydroxy-2-(1H-imidazol-1-yl)ethylidene-1,1-bisphosphonic acid (zolendronate).

A non-limiting class of bisphosphonate actives useful in the instant invention are selected from the group consisting of alendronate, cimadronate, clodronate, tiludronate, etidronate, ibandronate, neridronate, olpadronate, risedronate, piridronate, pamidronate, zolendronate, pharmaceutically acceptable salts thereof, and mixtures thereof.

A non-limiting subclass of the above-mentioned class in the instant case is selected from the group consisting of alendronate, pharmaceutically acceptable salts thereof, and mixtures thereof.

A non-limiting example of the subclass is alendronate monosodium trihydrate.

In the present invention, as it relates to bone stimulation, the agonist is typically administered for a sufficient period of time until the desired therapeutic effect is achieved. The term "until the desired therapeutic effect is achieved", as used herein, means that the therapeutic agent or agents are continuously administered, according to the dosing schedule chosen, up to the time that the clinical or medical effect sought for the disease or condition being mediated is observed by the clinician or researcher. For methods of treatment of the present invention, the compounds are continuously administered until the desired change in bone mass or structure is observed. In such instances, achieving an increase in bone mass or a replacement of abnormal bone structure with normal bone structure are the desired objectives. For methods of reducing the risk of a disease state or condition, the compounds are continuously administered for as long as necessary to prevent the undesired condition. In such instances, maintenance of bone mass density is often the objective.

Nonlimiting examples of administration periods can range from about 2 weeks to the remaining lifespan of the mammal. For humans, administration periods can range from about 2 weeks to the remaining lifespan of the human, preferably from about 2 weeks to about 20 years, more preferably from about 1 month to about 20 years, more preferably from about 6 months to about 10 years, and most preferably from about 1 year to about 10 years.

The instant compounds are also useful in combination with known agents useful for treating or preventing bone loss, bone fractures, osteoporosis, glucocorticoid induced osteoporosis, Paget's disease, abnormally increased bone turnover, periodontal disease, tooth loss, osteoarthritis, rheumatoid arthritis, , periprosthetic osteolysis, osteogenesis imperfecta, metastatic bone disease, hypercalcemia of malignancy, and multiple myeloma. Combinations of the presently disclosed compounds with other agents useful in treating or preventing osteoporosis or other bone disorders are within the scope of the invention. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the disease involved. Such agents include the following: an organic bisphosphonate; a cathepsin K inhibitor; an estrogen or an estrogen receptor modulator; an androgen receptor modulator; an inhibitor of osteoclast proton ATPase; an inhibitor of HMG-CoA reductase; an integrin receptor antagonist; an osteoblast anabolic agent, such as PTH; calcitonin; Vitamin D or a synthetic Vitamin D analogue; and the pharmaceutically acceptable salts and mixtures thereof. A preferred combination is a compound of the present invention and an organic bisphosphonate. Another preferred combination is a compound of the present invention and an estrogen receptor modulator. Another preferred combination is a compound of the present invention and an estrogen. Another preferred combination is a compound of the

present invention and an androgen receptor modulator. Another preferred combination is a compound of the present invention and an osteoblast anabolic agent.

Regarding treatment of abnormal bone resorption and ocular disorders, the formula I agonists generally have an EC<sub>50</sub> value from about 0.001 nM to about 100 microM,

5 although agonists with activities outside this range can be useful depending upon the dosage and route of administration. In a subclass of the present invention, the agonists have an EC<sub>50</sub> value of from about 0.01 microM to about 10 microM. In a further subclass of the present invention, the agonists have an EC<sub>50</sub> value of from about 0.1 microM to about 10 microM. EC<sub>50</sub> is a common measure of agonist activity well known to those of ordinary skill in the art and is  
10 defined as the concentration or dose of an agonist that is needed to produce half, i.e. 50%, of the maximal effect. See also, Goodman and Gilman's, *The Pharmacologic Basis of Therapeutics*, 9th edition, 1996, chapter 2, E. M. Ross, *Pharmacodynamics, Mechanisms of Drug Action and the Relationship Between Drug Concentration and Effect*, and PCT US99/23757, filed October 12, 1999, which are incorporated by reference herein in their entirety.

15 The herein examples illustrate but do not limit the claimed invention. Each of the claimed compounds are EP4 agonists and are useful for a number of physiological ocular and bone disorders.


Some abbreviations that may appear in this application are as follows:

20 ABBREVIATIONS

Designation

	CDI	1,1'-carbonyldiimidazole
	DHP	4-dihydro-2H-pyran
	LiOH	lithium hydroxide
25	NaBH <sub>4</sub>	sodium borohydride
	NaH	sodium hydride
	nBu <sub>3</sub> SnN <sub>3</sub>	azidotributyltin.
	PG	protecting groups
	TBSCl	tert-butyldimethylsilyl chloride
30	TsOH	p-toluenesulfonic acid

Compounds stated in the present invention can be prepared according to the following general scheme. All variables are as defined above unless otherwise indicated.





Preparation of compounds in the present invention is further illustrated by the following specific example. Following the general scheme described above and the procedure exemplified in the example, compounds of the invention have alternative groups for the defined variables can be prepared, e.g. following the procedure below, methyl 5-(3-bromopropyl)thiophene-2-carboxylate can be replaced with methyl 5-(3-bromopropyl)benzyl-2-carboxylate, or methyl 5-(3-bromopropyl)thiazole-2-carboxylate, to make the corresponding pyrrolidinone.

## EXAMPLE 1

5-(3-((2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)thiophene-2-carboxylic acid (9 and 10)

The preparation of compounds 9 and 10 was carried out according to the follow scheme.





The preparation of 1 was carried out according to the literature procedure (see: Tetrahedron 1994, 6221)

5 (5R)-5-[(1E)-4,4-difluoro-3-oxo-4-phenylbut-1-enyl]-1-(4-methoxybenzyl)pyrrolidin-2-one (2)

At -72 °C, oxalyl chloride (544 µL) was added dropwise to a solution of dimethylsulfoxide (480 µL) in CH<sub>2</sub>Cl<sub>2</sub> (14 ml) and the mixture was stirred 20 min at that temperature. A solution of (5R)-5-(hydroxymethyl)-1-(4-methoxybenzyl)pyrrolidin-2-one (714 mg, 3.04 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 ml) was then added slowly and the mixture was stirred for an hour at -72 °C. Triethylamine (2.0 ml) was then added dropwise and the mixture was allowed to warm to 0 °C. Water was added and the product was extracted in CH<sub>2</sub>Cl<sub>2</sub>, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to dryness. It was used as such in the next step. <sup>1</sup>H NMR (acetone-d<sub>6</sub>) δ 9.50 (s, 1H), 7.20 (d, 2H), 6.90 (d, 2H), 4.80 (d, 1H), 4.14 (d, 1H), 4.06 (m, 1H), 3.79, (s, 3H), 2.20-2.45 (m, 3H), 2.12 (m, 1H).

15

To a solution of dimethyl 3,3-difluoro-2-oxo-3-phenylpropylphosphonate (United States Patent 4,320,136 March. 16, 1982) (2.076 g) in THF (17 mL) at 0 °C was added potassium tert-butoxide (963 mg) and the mixture was stirred for an additional 1 hour at 0 °C. To the mixture was then added 2R)-1-(4-methoxybenzyl)-5-oxopyrrolidine-2-carbaldehyde in THF

(10 mL) via cannula and the resultant mixture stirred at room temperature 2 hours and quenched with saturated  $\text{NH}_4\text{Cl}$ . The mixture was then extracted with ethyl acetate (3x) and the organic layer was washed with water, brine, dried over  $\text{Mg}_2\text{SO}_4$ , filtered and concentrated. The residue was purified by chromatography using 20% acetone/toluene as the eluent to give the desired product 2.

5-(4,4-Difluoro-3-hydroxy-4-phenyl-but-1-enyl)-1-(4-methoxy-benzyl)-pyrrolidin-2-one (3)

To a solution of 2 (8.2 g, 21.2 mmol) in 80 mL  $\text{CH}_2\text{Cl}_2$  was added 1M (S)-CBS in toluene (10.6 mL, 10.6 mmol) and cooled to  $-40^\circ\text{C}$  to which a solution of catechol borane (6.8 mL, 63.8 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 mL) was added dropwise. The solution was stirred at  $-40^\circ\text{C}$  for one hour and allowed to warm up to  $-20^\circ\text{C}$  during the following two hours. The reaction mixture was quenched at  $-20^\circ\text{C}$  with 1 N HCl and was stirred for 4 hours at room temperature. The phases were separated and the organic phase was sequentially washed with 1N HCl,  $\text{H}_2\text{O}$ , 1 N NaOH, brine and dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo. The compound was purified by flash chromatography using 40-50% ethyl acetate/hexanes to give the desired product 3 as a pale yellow oil. MS ( $M + 1$ ) 388.2

(5*R*)-5-((1*E*)-3-([*tert*-butyl(dimethyl)silyl]oxy)-4,4-difluoro-4-phenylbut-1-enyl)-1-(4-methoxybenzyl)pyrrolidin-2-one (4)

To a solution 3 (365 mg) in DMF (3 mL) at room temperature was added imidazole (139 mg) followed by TBSCl (220 mg). The mixture was stirred over the weekend and then quenched with water. The mixture was extracted with ether (3x) and washed with water, brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo. Purified by column chromatography (50% ethyl acetate : hexane) afforded compound 4.

(5*R*)-5-((1*E*)-3-([*tert*-butyl(dimethyl)silyl]oxy)-4,4-difluoro-4-phenylbut-1-enyl)pyrrolidin-2-one (5)

To a solution of 4 (359 mg) in acetonitrile (20 mL) at  $0^\circ\text{C}$  was added CAN (2 g), water (2 mL) and the mixture was allowed to warm to room temperature for 4 hours. The mixture was extracted with ether (3x) and was washed with water, brine and dried over  $\text{Na}_2\text{SO}_4$ . Purification by column chromatography (50%-75%-100% ethyl acetate in hexane) afforded compound 5.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.50-7.40 (m, 5H), 5.70-5.65 (m, 2H), 4.50-4.42 (m, 1H), 4.20-4.13 (m, 1H), 2.37-2.30 (m, 3H), 1.80-1.70 (m, 1H), 0.87 (s, 9H), -0.05 (d, 6H).

methyl 5-{3-[(2R)-2-((1E)-3-{[*tert*-butyl(dimethyl)silyl]oxy}-4,4-difluoro-4-phenylbut-1-enyl)-5-oxopyrrolidin-1-yl]propyl}thiophene-2-carboxylate (6)

To a solution of NaH 60% (30 mg) in DMF (2 mL) was added 5 (182 mg) in DMF (2mL), methyl 5-(3-bromopropyl)thiophene-2-carboxylate (200 mg) in DMF (1.5 mL) and NaI (30 mg). The mixture was stirred at 50 °C for 3h. After cooling to room temperature, the mixture was quenched with saturated NH<sub>4</sub>Cl and extracted with ethyl acetate (3 x). The organic layer was washed with water, brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The crude was purified by flash chromatography. Eluting with 50% ethyl acetate in hexanes gave the desired product 6.

10 methyl 5-(3-{(2R)-2-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}propyl)thiophene-2-carboxylate (7 and 8)

To a solution of 6 (230 mg) in THF (5 mL) was added TBAF (1.0 M in THF, 0.6 mL) and the mixture was stirred at room temperature for 30 min. The reaction mixture was concentrated in vacuo and purified by column chromatography (ethyl acetate) affording a mixture of compounds 7 (less polar) and 8 (more polar). The isomers were separated by HPLC (Chiralpak AD<sup>®</sup>) using 30% isopropanol in hexanes. Isomer 7 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.65 (d, 1H), 7.49-7.42 (m, 5H), 6.83 (d, 1H), 5.72-5.60 (m, 2H), 4.61-4.55 (m, 1H), 4.08-4.02 (m, 1H), 3.88 (s, 3H), 3.54-3.46 (m, 1H), 2.89-2.78 (m, 3H), 2.40-2.32 (m, 2H), 2.21-2.14 (m, 1H), 1.87-1.77 (m, 2H), 1.70-1.63 (m, 1H). Isomer 8 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.65 (d, 1H), 7.49-7.43 (m, 5H), 6.83 (d, 1H), 5.72-5.61 (m, 2H), 4.61-4.54 (m, 1H), 4.07-4.02 (m, 1H), 3.88 (s, 3H), 3.54-3.47 (m, 1H), 2.87-2.79 (m, 3H), 2.44-2.28 (m, 2H), 2.22-2.13 (m, 1H), 1.89-1.76 (m, 2H), 1.72-1.64 (m, 1H).

The title compounds: 5-(3-{(2R)-2-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}propyl)thiophene-2-carboxylic acid (9 and 10)

A mixture of ester 7 or 8 in methanol (4.7 mL), water (1 mL) and LiOH (0.5mL, 1.0M) was stirred at room temperature under N<sub>2</sub> overnight and concentrated to give the title compound as a lithium salt. The salt was washed with ether, acidified with HCl (1.0 N), and extracted with ethyl acetate (3x). The extracts were washed with water, brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give the title compound 9 (less polar) or 10 (more polar). MS (-ESI): m/z 434.1 (M-1).



### Effects of an EP4 Agonist on Intraocular Pressure (IOP) in Rabbits and Monkeys

Animals - Drug-naïve, male Dutch Belted rabbits and female cynomolgus monkeys are used in this study. Animal care and treatment in this investigation are in compliance with guidelines by the National Institute of Health and the Association for Research in Vision and Ophthalmology resolution in the use of animals for research. All experimental procedures str approved by the Institutional Animal Care and Use Committee of Merck and Company.

Drug Preparation and Administration - Drug concentrations are expressed in terms of the active ingredient (base). The compounds of this invention are dissolved in physiological saline at 0.01, 0.001, 0.0001 % for rabbit study and 0.05, 0.005% for monkey studies. Drug or vehicle aliquots (25 ul) are administered topically unilaterally or bilaterally. In unilateral applications, the contralateral eyes receive an equal volume of saline. Proparacaine (0.5%) is applied to the cornea prior to tonometry to minimize discomfort. Intraocular pressure (IOP) is recorded using a pneumatic tonometer (Alcon Applanation Pneumatograph) or equivalent.

Statistical Analysis - The results are expressed as the changes in IOP from the basal level measured just prior to administration of drug or vehicle and represent the mean, plus or minus standard deviation. Statistical comparisons are made using the Student's t-test for non-paired data between responses of drug-treated and vehicle-treated animals and for paired data between ipsilateral and contralateral eyes at comparable time intervals. The significance of the date is also determined as the difference from the "t-0" value using Dunnett's "t" test. Asterisks represent a significance level of  $p < 0.05$ .

Intraocular Pressure Measurement in Rabbits - Male Dutch Belted rabbits weighing 2.5-4.0 kg are maintained on a 12- hour light/dark cycle and rabbit chow. All experiments are performed at the same time of day to minimize variability related to diurnal rhythm. IOP is measured before treatment then the compounds of this invention or vehicle are instilled (one drop of 25 ul) into one or both eyes and IOP is measured at 30, 60, 120, 180, 240, 300, and 360 minutes after instillation. In some cases, equal number of animals treated bilaterally with vehicle only are evaluated and compared to drug treated animals as parallel controls.

Intraocular Pressure Measurements in Monkeys - Unilateral ocular hypertension of the right eye is induced in female cynomolgus monkeys weighing between 2 and 3 kg by photocoagulation of the trabecular meshwork with an argon laser system (Coherent NOVUS 2000, Palo Alto, USA)

using the method of Lee et al. (1985). The prolonged increase in intraocular pressure (IOP) results in changes to the optic nerve head that are similar to those found in glaucoma patients. For IOP measurements, the monkeys are kept in a sitting position in restraint chairs for the duration of the experiment. Animals are lightly anesthetized by the intramuscular injection of ketamine hydrochloride (3-5 mg/kg) approximately five minutes before each IOP measurement and one drop of 0.5% proparacaine was instilled prior to recording IOP. IOP is measured using a pneumatic tonometer (Alcon Applanation Tonometer) or a Digilab pneumatonometer (Bio-Rad Ophthalmic Division, Cambridge, MA, USA). IOP is measured before treatment and generally at 30, 60, 124, 180, 300, and 360 minutes after treatment. Baseline values are also obtained at these time points generally two or three days prior to treatment. Treatment consists of instilling one drop of 25  $\mu$ l of the compounds of this invention (0.05 and 0.005 %) or vehicle (saline). At least one-week washout period is employed before testing on the same animal. The normotensive (contralateral to the hypertensive) eye is treated in an exactly similar manner to the hypertensive eye. IOP measurements for both eyes are compared to the corresponding baseline values at the same time point. Results are expressed as mean plus-or-minus standard deviation in mm Hg. The activity range of the compounds of this invention for ocular use is between 0.01 and 100,000 nM

Radioligand binding assays - The assays used to test these compounds were performed essentially as described in: Abramovitz M, Adam M, Boie Y, Carriere M, Denis D, Godbout C, Lamontagne S, Rochette C, Sawyer N, Tremblay NM, Belley M, Gallant M, Dufresne C, Gareau Y, Ruel R, Juteau H, Labelle M, Ouimet N, Metters KM. The utilization of recombinant prostanoid receptors to determine the affinities and selectivities of prostaglandins and related analogs. *Biochim Biophys Acta* 2000 Jan 17;1483(2):285-293 and discussed below:

Stable expression of prostanoid receptors in the human embryonic kidney (HEK) 293(EBNA) cell line - Prostanoid receptor (PG) cDNAs corresponding to full length coding sequences were subcloned into the appropriate sites of the mammalian expression vector pCEP4 (Invitrogen) pCEP4PG plasmid DNA was prepared using the Qiagen plasmid preparation kit (QIAGEN) and transfected into HEK 293(EBNA) cells using LipofectAMINE<sup>®</sup> (GIBCO-BRL) according to the manufacturers' instructions. HEK 293(EBNA) cells expressing the cDNA together with the hygromycin resistance gene were selected in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10 % heat inactivated fetal bovine serum, 1 mM sodium pyruvate, 100 U/ml Penicillin-G, 100  $\mu$ g/ml Streptomycin sulphate, 250  $\mu$ g/ml active GENETICIN<sup>™</sup> (G418) (all from Life Technologies, Inc./BRL) and 200  $\mu$ g/ml hygromycin (Calbiochem). Individual

colonies were isolated after 2-3 weeks of growth under selection using the cloning ring method and subsequently expanded into clonal cell lines. Expression of the receptor cDNA was assessed by receptor binding assays. HEK 293(EBNA) cells were grown in supplemented DMEM complete medium at 37°C in a humidified atmosphere of 6 % CO<sub>2</sub> in air, then harvested and membranes prepared by differential centrifugation (1000 x g for 10 min, then 160,000 x g for 30 min, all at 4°C) following lysis of the cells by nitrogen cavitation at 800 psi for 30 min on ice in the presence of protease inhibitors (2 mM phenylmethylsulfonylfluoride, 10 µM E-64, 100 µM leupeptin and 0.05 mg/ml pepstatin). The 160,000 x g pellets were resuspended in 10 mM HEPES/KOH (pH 7.4) containing 1 mM EDTA at approximately 5-10 mg/ml protein by Dounce homogenisation (Dounce A; 10 strokes), frozen in liquid nitrogen and stored at -80°C.

Prostanoid receptor binding assays - Prostanoid receptor binding assays were performed in a final incubation volume of 0.2 ml in 10 mM MES/KOH (pH 6.0) (EP subtypes, FP and TP) or 10 mM HEPES/KOH (pH 7.4) (DP and IP), containing 1 mM EDTA, 10 mM MgCl<sub>2</sub> (EP subtypes) or 10 mM MnCl<sub>2</sub> (DP, FP, IP and TP) and radioligand [0.5-1.0 nM [<sup>3</sup>H]PGE<sub>2</sub> (181 Ci/mmol) for EP subtypes, 0.7 nM [<sup>3</sup>H]PGD<sub>2</sub> (115 Ci/mmol) for DP, 0.95 nM [<sup>3</sup>H]PGF<sub>2α</sub> (170 Ci/mmol) for FP, 5 nM [<sup>3</sup>H]iloprost (16 Ci/mmol) for IP and 1.8 nM [<sup>3</sup>H]SQ 29548 (46 Ci/mmol) for TP]. EP<sub>3</sub> assays also contained 100 µM GTPγS. The reaction was initiated by addition of membrane protein (approximately 30 µg for EP<sub>1</sub>, 20 µg for EP<sub>2</sub>, 2 µg for EP<sub>3</sub>, 10 µg for EP<sub>4</sub>, 60 µg for FP, 30 µg for DP, 10 µg for IP and 10 µg for TP) from the 160,000 x g fraction. Ligands were added in dimethylsulfoxide (Me<sub>2</sub>SO) which was kept constant at 1 % (v/v) in all incubations. Non-specific binding was determined in the presence of 1 µM of the corresponding non-radioactive prostanoid. Incubations were conducted for 60 min (EP subtypes, FP and IP) or 30 min (DP and TP) at 30°C (EP subtypes, DP, FP and TP) or room temperature (IP) and terminated by rapid filtration through a 96-well Unifilter GF/C (Canberra Packard) prewetted in assay incubation buffer without EDTA (at 4°C) and using a Tomtec Mach III 96-well semi-automated cell harvester. The filters were washed with 3-4 ml of the same buffer, dried for 90 min at 55°C and the residual radioactivity bound to the individual filters determined by scintillation counting with addition of 50 µl of Ultima Gold F (Canberra Packard) using a 1450 MicroBeta (Wallac). Specific binding was calculated by subtracting non-specific binding from total binding. Specific binding represented 90-95 % of the total binding and was linear with respect to the concentrations of radioligand and protein used. Total binding represented 5-10 % of the radioligand added to the incubation media. The activity range of the compounds of this invention for bone use is between 0.01 and 100,000 nM.

### Bone Resorption Assays

Animal Procedures - For mRNA localization experiments, 5-week old Sprague-Dawley rats (Charles River) are euthanized by CO<sub>2</sub>, their tibiae and calvariae are excised, cleaned of soft tissues and frozen immediately in liquid nitrogen. For EP<sub>4</sub> regulation experiments, 6-week old

5 rats are given a single injection of either vehicle (7% ethanol in sterile water) or an anabolic dose of PGE<sub>2</sub> (Cayman Chemical, Ann Arbor, MI), 3-6 mg/kg in the same vehicle) intraperitoneally.

Animals are euthanized at several time points post-injection and their tibiae and calvariae, as well as samples from lung and kidney tissues are frozen in liquid nitrogen.

10 Cell Cultures - RP-1 periosteal cells are spontaneously immortalized from primary cultures of periosteal cells from tibiae of 4-week old Sprague-Dawley rats and are cultured in DMEM (BRL, Gaithersburg, MD) with 10 % fetal bovine serum (JRH Biosciences, Lenexa, KS). These cells do not express osteoblastic phenotypic markers in early culture, but upon confluence, express type I collagen, alkaline phosphatase and osteocalcin and produce mineralized extracellular

15 matrix. RCT-1 and RCT-3 are clonal cell lines immortalized by SV-40 large T antigen from cells released from fetal rat calvaria by a combination collagenase/hyaluronidase digestion. RCT-1 cells, derived from cells released during the first 10 minutes of digestion (fraction I), are cultured in RPMI 1640 medium (BRL) with 10% fetal bovine serum and 0.4 mg/ml G418 (BRL). These cells differentiate and express osteoblastic features upon retinoic acid treatment. RCT-3 cells,

20 immortalized from osteoblast-enriched fraction III cells, are cultured in F-12 medium (BRL) with 5% Fetal bovine serum and 0.4 mg/ml G418. TRAB-11 cells are also immortalized by SV40 large T antigen from adult rat tibia and are cultured in RPMI 1640 medium with 10% FBS and 0.4 mg/ml G418. ROS 17/2.8 rat osteosarcoma cells are cultured in F-12 containing 5% FBS. Osteoblast-enriched (fraction III) primary fetal rat calvaria cells are obtained by

25 collagenase/hyaluronidase digestion of calvariae of 19 day-old rat fetuses. See Rodan et al., *Growth stimulation of rat calvaria osteoblastic cells by acidic FGF*, *Endocrinology*, 121, 1919-1923 (1987), which is incorporated by reference herein in its entirety. Cells are released during 30-50 minutes digestion (fraction III) and are cultured in F-12 medium containing 5% FBS. P815 (mouse mastocytoma) cells, cultured in Eagles MEM with 10% FBS, and NRK (normal rat

30 kidney fibroblasts) cells, cultured in DMEM with 10% FBS, are used as positive and negative controls for the expression of EP<sub>4</sub>, respectively. See Abramovitz et al., Human prostanoid receptors: cloning and characterization. In: Samulesson B. et al. ed) *Advances in prostaglandin, Thromboxanes and leukotriene research*, vol. 23, pp. 499-504 (1995) and de Larco et al., Epithelioid and fibroblastic rat kidney cell clones: EGF receptors and the effect of mouse

sarcoma virus transformation, *Cell Physiol.*, 94, 335-342 (1978), which are both incorporated by reference herein in their entirety.

Northern Blot Analysis - Total RNA is extracted from the tibial metaphysis or diaphysis and calvaria using a guanidinium isothiocyanate-phenol-chloroform method after pulverizing frozen bone samples by a tissue homogenizer. See P. Chomczynski et al., Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction., *Analyt Biochem*, 162, 156-159 (1987), which is incorporated by reference herein in its entirety. RNA samples (20 mg) are separated on 0.9% agarose/formaldehyde gels and transferred onto nylon membranes (Boehringer Mannheim, Germany). Membranes are prehybridized in Hybrisol I (Oncor, Gaithersburg, MD) and 0.5 mg/ml sonicated salmon sperm DNA (Boehringer) at 42°C for 3 hours and are hybridized at 42°C with rat EP<sub>2</sub> and mouse EP<sub>4</sub> cDNA probes labeled with [<sup>32</sup>P]-dCTP (Amersham, Buckinghamshire, UK) by random priming using the rediprime kit (Amersham). After hybridization, membranes are washed 4 times in 2xSSC + 0.1% SDS at room temperature for a total of 1 hour and once with 0.2xSSC + 0.1% SDS at 55°C for 1 hour and then exposed to Kodak XAR 2 film at -70°C using intensifying screens. After developing the films, bound probes are removed twice with 0.1% SDS at 80°C and membranes are hybridized with a human GAPDH (Glyceraldehyde 3-Phosphate Dehydrogenase) cDNA probe (purchased from Clontech, Palo Alto, CA) for loading control.

In-Situ Hybridization - Frozen tibiae are sectioned coronally at 7 mm thickness and sections are mounted on charged slides (Probe On Plus, Fisher Scientific, Springfield, NJ) and are kept at -70°C until hybridization. cRNA probes are labeled with <sup>35</sup>S-UTPgS (ICN, Costa Mesa, CA) using a Riboprobe II kit (Promega Madison, WI). Hybridization is performed overnight at 50° C. See M. Weinreb et al., *Different pattern of alkaline phosphatase, osteopontin and osteocalcin expression in developing rat bone visualized by in-situ hybridization*, *J. Bone Miner Res.*, 5, 831-842 (1990) and D. Shinar et al., *Expression of alphav and beta3 integrin subunits in rat osteoclasts in situ*, *J. Bone Miner. Res.*, 8, 403-414 (1993), which are both incorporated by reference herein in their entirety. Following hybridization and washing, sections are dipped in Ilford K5 emulsion diluted 2:1 with 6% glycerol in water at 42° C and exposed in darkness at 4° C for 12-14 days. Slides are developed in Kodak D-19 diluted 1:1 with water at 15°, fixed, washed in distilled water and mounted with glycerol-gelatin (Sigma) after hematoxylin staining. Stained sections are viewed under the microscope (Olympus, Hamburg, Germany), using either bright-field or dark-field optics.

- Expression Of EP<sub>4</sub> In Osteoblastic Cell Lines And In Bone Tissue - The expression of EP<sub>4</sub> and EP<sub>2</sub> mRNA is examined in various bone derived cells including osteoblast-enriched primary rat calvaria cells, immortalized osteoblastic cell lines from fetal rat calvaria or from adult rat tibia and an osteoblastic osteosarcoma cell line. Most of the osteoblastic cells and cell lines show significant amounts of 3.8 kb EP<sub>4</sub> mRNA, except for the rat osteosarcoma cell line ROS 17/2.8. Consistent with this finding, in ROS 17/2.8 cells PGE<sub>2</sub> has no effect on intracellular cAMP, which is markedly induced in RCT-3 and TRAB-11 cells. Treatment of RCT-1 cells with retinoic acid, which promotes their differentiation, reduces the levels of EP<sub>4</sub> mRNA. NRK fibroblasts do not express EP<sub>4</sub> mRNA, while P815 mastocytoma cells, used as positive controls, express large amounts of EP<sub>4</sub> mRNA. In contrast to EP<sub>4</sub> mRNA, none of the osteoblastic cells and cell lines express detectable amounts of EP<sub>2</sub> mRNA in total RNA samples. Expression of EP<sub>4</sub> mRNA in osteoblastic cells, EP<sub>4</sub> is also expressed in total RNA isolated from tibiae and calvariae of 5-week-old rats. In contrast, no EP<sub>2</sub> mRNA is found in RNA from tibial shafts.
- 15 PGE<sub>2</sub> Induces The Expression Of EP<sub>4</sub> mRNA in RP-1 Periosteal Cells And In Adult Rat Tibiae - PGE<sub>2</sub> enhances its own production via upregulation of cyclooxygenase 2 expression in osteoblasts and in bone tissue thus autoamplifying its own effects. PGE<sub>2</sub> also increases the levels of EP<sub>4</sub> mRNA. RP-1 cells are immortalized from a primary culture of adult rat tibia periosteum is examined. These cells express osteoblast phenotypic markers upon confluence and form mineralized bone matrix when implanted in nude mice. Similar to the other osteoblastic cells examined, RP-1 periosteal cells express a 3.8 kb EP<sub>4</sub> transcript. Treatment with PGE<sub>2</sub> (10<sup>-6</sup> M) rapidly increases EP<sub>4</sub> mRNA levels peaking at 2 hours after treatment. PGE<sub>2</sub> has no effect on EP<sub>4</sub> mRNA levels in the more differentiated RCT-3 cells pointing to cell-type specific regulation of EP<sub>4</sub> expression by PGE<sub>2</sub>. EP<sub>2</sub> mRNA is not expressed in RP-1 cells before or after treatment with PGE<sub>2</sub>. To examine if PGE<sub>2</sub> regulates EP<sub>4</sub> mRNA levels *in vivo* in bone tissue, five-week-old male rats are injected with PGE<sub>2</sub> (3 - 6 mg/Kg). Systemic administration of PGE<sub>2</sub> rapidly increased EP<sub>4</sub> mRNA levels in the tibial diaphysis peaking at 2 h after injection. A similar effect of PGE<sub>2</sub> on EP<sub>4</sub> mRNA is observed in the tibial metaphysis and in calvaria. PGE<sub>2</sub> induces EP<sub>4</sub> mRNA levels *in vitro* in osteogenic periosteal cells and *in vivo* in bone tissue in a cell type-specific and tissue-specific manner. PGE<sub>2</sub> does not induce EP<sub>2</sub> mRNA in RP-1 cells nor in bone tissue.

Localization of EP<sub>4</sub> mRNA expression in bone tissue - *In situ* hybridization is used in order to localize cells expressing EP<sub>4</sub> in bone. In control experiment (vehicle-injected) rats, low


expression of EP<sub>4</sub> is detected in bone marrow cells. Administration of a single anabolic dose of PGE<sub>2</sub> increased the expression of EP<sub>4</sub> in bone marrow cells. The distribution of silver grains over the bone marrow is not uniform and occurs in clumps or patches in many areas of the metaphysis. Within the tibial metaphysis, EP<sub>4</sub> expression is restricted to the secondary

5 spongiosa area and is not seen in the primary spongiosa. Hybridization of similar sections with a sense probe (negative control) does not show any signal. EP<sub>4</sub> is expressed in osteoblastic cells *in vitro* and in bone marrow cells *in vivo*, and is upregulated by its ligand, PGE<sub>2</sub>.

10 Agonist activity - Using standard methods for measuring agonist activity, the compounds of the invention were evaluated in cell cultures and in EP<sub>4</sub> receptor cell-free systems to determine the agonist activity of the compounds in terms of their EC<sub>50</sub> value.

WHAT IS CLAIMED IS:

1. A compound having the structural formula I:




FORMULA I

or a pharmaceutically acceptable salt thereof, wherein,

Y¹ is

1) CH<sub>2</sub>CH<sub>2</sub>,

2) CHCH, or



Y is C(O) or CH(OH);

A is (CH<sub>2</sub>)<sub>n</sub>;


n is 1, 2, 3, or 4;

W a bond, unsubstituted C 1-6 alkylene, or C 1-6 alkylene substituted with 1, 2, 3, or 4 halogen atoms;

Z is

1) O,

2) S,




4) HC=CH,

5) C≡C, or


6) a bond;

Q is a disubstituted aryl or heteroaryl ring, wherein one ring atom of the ring is attached to the moiety



and another ring atom is attached to the moiety





R<sup>1</sup> is

COR<sup>5</sup>,

OH,

CN,

(CH<sub>2</sub>)<sub>1-3</sub> CO<sub>2</sub>R<sup>6</sup>,

C(O)NHSO<sub>2</sub>R<sup>8</sup>,

SO<sub>2</sub>R<sup>7</sup>,

(CH<sub>2</sub>)<sub>0-4</sub>SO<sub>3</sub>R<sup>6</sup>,

CF<sub>2</sub>SO<sub>2</sub>NH<sub>2</sub>,

SO<sub>2</sub>NH<sub>2</sub>,

SO<sub>2</sub>NHCOR<sup>8</sup>,

PO(OR<sup>7</sup>)<sub>2</sub>,

C<sub>1-4</sub> alkoxy,

hydroxymethylketone, or

(CH<sub>2</sub>)<sub>0-4</sub>R<sup>k</sup>, wherein R<sup>k</sup> is unsubstituted or substituted with 1 to 3 groups of R<sup>a</sup>;

R<sup>2</sup> is

1) C<sub>1-6</sub>alkyl,

2) (CH<sub>2</sub>)<sub>0-8</sub>C<sub>6-10</sub>aryl,

3) (CH<sub>2</sub>)<sub>0-8</sub>R<sup>m</sup>,

4) (CH<sub>2</sub>)<sub>0-8</sub>C<sub>3-8</sub>cycloalkyl,

5) O-C<sub>1-10</sub>alkyl,

6) O-C<sub>6-10</sub>aryl,

7) O-R<sup>m</sup>,

8) O-C<sub>3-10</sub>cycloalkyl

wherein aryl, R<sup>m</sup>, and cycloalkyl are unsubstituted or substituted with 1-3 groups of R<sup>b</sup>;

R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of

1) halogen, and

2) C<sub>1-6</sub> alkyl, or

R<sup>3</sup> and R<sup>4</sup>, together with the carbon atom to which they are attached, form a C<sub>3-7</sub> cycloalkyl ring;

R<sup>5</sup> is

- 1) hydrogen,
- 2) OH,
- 3) CH<sub>2</sub>OH,
- 5 4) C<sub>1-6</sub> alkoxy,
- 5) NHPO<sub>2</sub>R<sup>6</sup>,
- 6) NHR<sup>9</sup>,
- 7) NHSO<sub>2</sub>R<sup>8</sup>, or
- 8) NR<sup>6</sup>R<sup>7</sup>;

- 10 R<sup>6</sup> and R<sup>7</sup> are independently selected from the group consisting of hydrogen, C<sub>1-6</sub> alkyl, and C<sub>3-8</sub> cycloalkyl;

R<sup>8</sup> is selected from the group consisting of hydrogen, C<sub>6-10</sub>aryl, R<sup>n</sup>, and C<sub>1-4</sub>alkyl;

R<sup>9</sup> is C(O)R<sup>10</sup> or SO<sub>2</sub>R<sup>10</sup>;

R<sup>10</sup> is hydrogen, C<sub>6-10</sub> aryl, or C<sub>1-4</sub> alkyl;

- 15 R<sup>a</sup> and R<sup>b</sup> are independently selected from the group consisting of

- 1) C<sub>1-6</sub>alkoxy,
- 2) C<sub>1-6</sub>alkyl, unsubstituted or substituted with
  - a) C<sub>1-6</sub> alkoxy,
  - b) C<sub>1-6</sub> alkylthio,
- 20 c) CN,
- d) OH, or
- e) CF<sub>3</sub>,

3) CF<sub>3</sub>,

4) nitro,

- 25 5) amino,

6) cyano,

7) C<sub>1-6</sub>alkylamino,

8) halogen

9) OR<sup>c</sup>,

- 30 10) OCH<sub>2</sub>R<sup>c</sup>, and

11) CH<sub>2</sub>OR<sup>c</sup>;

R<sup>c</sup> is

1) C<sub>6-10</sub>aryl,

2) R<sup>s</sup>, or

- 35 3) C<sub>3-8</sub>cycloalkyl; and

$R^k$ ,  $R^m$ ,  $R^n$  and  $R^s$  are independently selected from the group consisting of


- 1) a stable monocyclic heteroaryl ring having 5, 6 or 7 ring atoms, or a stable bicyclic heteroaryl ring having 8, 9, 10, or 11 ring atoms, wherein the monocyclic ring has 1, 2, 3, or 4 heteroatoms, independently selected from the group consisting of O, S or N, and wherein the bicyclic ring has 1, 2, 3, or 4 heteroatoms, independently selected from the group consisting of O, S or N, and
- 2) a stable monocyclic or bicyclic heterocycloalkyl ring system a stable, saturated monocyclic or bicyclic ring system having 3 to 10 ring atoms, wherein 1, 2, 3, or 4 ring atoms are heteroatoms selected from O, S and N.

2. The compound of Claim 1, or a pharmaceutically acceptable salt thereof, wherein  $Y^1$  is CHCH and Y is CH(OH).


3. The compound of Claim 2, or a pharmaceutically acceptable salt thereof, wherein A is  $(CH_2)_{1-3}$  and W is a bond or  $(CH_2)_{1-3}$ .

4. The compound of Claim 3, or a pharmaceutically acceptable salt thereof, wherein 1)  $R^1$  is COOH or tetrazole, 2)  $R^2$  is phenyl, and 3)  $R^3$  and  $R^4$  are independently selected from the group consisting of hydrogen and halogen, or  $R^3$  and  $R^4$  together with the carbon to which they are attached, form a cyclopropyl ring.

5. The compound of Claim 4, or a pharmaceutically acceptable salt thereof, wherein Q is selected from the group consisting of



6. The compound of Claim 5, or a pharmaceutically acceptable salt thereof, wherein Q is selected from the group consisting of



7. The compound of Claim 6 selected from the group consisting of

- (1) 5-(3-((2R)-2-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)thiophene-2-carboxylic acid,
- (2) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[5-(1H-tetrazol-5-yl)thien-2-yl]propyl}pyrrolidin-2-one,
- (3) 5-(3-((2R)-2-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)-1,3-thiazole-2-carboxylic acid,
- (4) 2-(3-((2R)-2-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)-1,3-thiazole-5-carboxylic acid,
- (5) (5R)-5-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[5-(1H-tetrazol-5-yl)-1,3-thiazol-2-yl]propyl}pyrrolidin-2-one,
- (6) 2-(3-((2R)-2-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)propyl)-1,3-thiazole-4-carboxylic acid,
- (7) [5-(2-((2R)-2-[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)ethyl)thien-2-yl]acetic acid,

(8) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{2-[5-(1*H*-tetraazol-5-ylmethyl)thien-2-yl]ethyl}pyrrolidin-2-one,

(9) 2-(3-[(2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]propyl)-1,3-oxazole-5-carboxylic acid,

(10) 5-(3-[(2*R*)-2-[(1*E*)-3-hydroxy-3-(1-phenylcyclopropyl)prop-1-enyl]-5-oxopyrrolidin-1-yl]propyl)thiophene-2-carboxylic acid,

(11) (5*R*)-5-[(1*E*)-3-hydroxy-3-(1-phenylcyclopropyl)prop-1-enyl]-1-{3-[5-(1*H*-tetraazol-5-yl)thien-2-yl]propyl}pyrrolidin-2-one,

(12) 5-(3-[(2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]propyl)-1,3,4-thiadiazole-2-carboxylic acid,

(13) 4-(3-[(2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]propyl)benzoic acid,

(14) 3-(3-[(2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]propyl)benzoic acid,

(15) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[3-(1*H*-tetraazol-5-yl)phenyl]propyl}pyrrolidin-2-one,

(16) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[4-(1*H*-tetraazol-5-yl)phenyl]propyl}pyrrolidin-2-one,

(17) 3-[5-[(2*R*)-2-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl)methyl]thien-2-yl]propanoic acid,

(18) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-({5-[2-(1*H*-tetraazol-5-yl)ethyl]thien-2-yl)methyl}pyrrolidin-2-one,

(19) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-({4-[3-(1*H*-tetraazol-5-yl)propyl]thien-3-yl)methyl}pyrrolidin-2-one,

(20) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-({4-[2-(1*H*-tetraazol-5-yl)ethyl]thien-2-yl)methyl}pyrrolidin-2-one,

(21) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-({4-[2-(1*H*-tetraazol-5-yl)ethyl]-1,3-thiazol-2-yl)methyl}pyrrolidin-2-one, and

(22) (5*R*)-5-[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-{3-[2-(1*H*-tetraazol-5-yl)ethyl]benzyl}pyrrolidin-2-one,

and pharmaceutically acceptable salts thereof.

8. A method for treating disorders related to elevated intraocular pressure by: treating ocular hypertension, treating glaucoma, treating macular edema, treating macular degeneration, increasing retinal and optic nerve head blood velocity, increasing retinal and optic nerve tension, providing a neuroprotective effect or treating dry eyes, comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof.

9. A topical composition comprising the compound of formula I as defined in any one of Claims 1 to 7, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

10. The composition of Claim 9, wherein the composition comprises xanthan gum or gellan gum.

11. The composition of Claim 10, wherein the composition is a solution or a suspension.

12. The method according to Claim 8 further comprising administering to the patient an active ingredient selected from the group consisting of a  $\beta$ -adrenergic blocking agent, a parasympatho-mimetic agent, a Maxi-K channel blocker, a sympathomimetic agent, a carbonic anhydrase inhibitor, a prostaglandin, a hypotensive lipid, a neuroprotectant, and a 5-HT<sub>2</sub> receptor agonist, is added to the formulation.

13. The method according to Claim 12 wherein the  $\beta$ -adrenergic blocking agent is timolol, betaxolol, levobetaxolol, carteolol, or levobunolol; the parasympathomimetic agent is pilocarpine; the Maxi-K channel blocker is Penitrem A, paspalicine, charybdotoxin, or iberiotoxin, the sympathomimetic agent is epinephrine, brimonidine, iopidine, clonidine, or para-aminoclonidine; the carbonic anhydrase inhibitor is dorzolamide, acetazolamide, metazolamide or brinzolamide; the prostaglandin is latanoprost, travaprost, unoprostone, rescala, or S1033; the hypotensive lipid is lumigan; the neuroprotectant is eliprodil, R-eliprodil or memantine; and the 5-HT<sub>2</sub> receptor agonist is 1-(2-aminopropyl)-3-methyl-1H-imidazol-6-ol fumarate or 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-ethylamine.

14. A compound of formula I of any one of Claims 1 to 7, or a pharmaceutically acceptable salt thereof, for use in medicinal therapy.

15. Use of a compound of formula I of any one of Claims 1 to 7, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating disorders related to elevated intraocular pressure.

16. Use of a compound of formula I of any one of Claims 1 to 7, or a pharmaceutically acceptable salt thereof, as a selective EP<sub>4</sub> receptor agonist.

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/CA 03/01620

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07D403/06

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; MARUYAMA, TORU ET AL: "Remedies for diseases with bone mass loss having EP4 agonist as the activ ingredient" retrieved from STN Database accession no. 138:142493 XP002277059 abstract &amp; WO 2003/009872 A (ONO PHARMACEUTICAL CO., LTD., JAPAN) 6 February 2003 (2003-02-06)</p> <p style="text-align: center;">--- -/--</p>	1,8,14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

**\* Special categories of cited documents :**

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

6 May 2004

Date of mailing of the international search report

03/06/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fanni, S



## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/CA 01/01620

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DATABASE WPI Section Ch, Week 200247 Derwent Publications Ltd., London, GB; Class B03, AN 2002-443972 XP002252111 abstract & WO 02/24647 A (ONO PHARM CO LTD) 28 March 2002 (2002-03-28) cited in the application ---	1,8,14
Y	EP 1 114 816 A (ONO PHARMACEUTICAL CO) 11 July 2001 (2001-07-11) page 6-7, definition of A claim 1 ---	1,8,14
Y,P	WO 03/008377 A (HOFFMANN LA ROCHE) 30 January 2003 (2003-01-30) claim 1 ---	1,8,14
E	WO 03/103772 A (MERCK FROSST CANADA INC ;BILLOT XAVIER (CA); HAN YONGXIN (CA); YOU) 18 December 2003 (2003-12-18) claim 1 examples 1,7 ---	1,8,14
E	WO 03/103604 A (APPLIED RESEARCH SYSTEMS ;BAO BAGNA (US); MCKENNA SEAN D (US); RED) 18 December 2003 (2003-12-18) claim 1 examples 5-15,17-28,37,38,43-50,56-59,65 -----	1,8,14

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/CA 03/01620

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 8,12,13,16 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

---

Continuation of Box I.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Information Publication No

PCT/CA 03/01620

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2003009872	A	06-02-2003	CA 2454584 A1 WO 03009872 A1	06-02-2003 06-02-2003
EP 1114816	A	11-07-2001	EP 1114816 A1 US 6586468 B1 WO 0015608 A1	11-07-2001 01-07-2003 23-03-2000
WO 03008377	A	30-01-2003	CA 2451392 A1 WO 03008377 A1 EP 1409455 A1 US 2003120079 A1	30-01-2003 30-01-2003 21-04-2004 26-06-2003
WO 03103772	A	18-12-2003	WO 03103772 A1	18-12-2003
WO 03103604	A	18-12-2003	WO 03103604 A2	18-12-2003

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**This Page Blank (uspto)**